

## A NOTE ON THE FUZZY XOR

RADOSŁAW ŁUKASIK 

**Abstract.** In this paper we will try to answer what conditions must be met by the fuzzy Xor to be used for cryptographic purposes. We will also show that defining the fuzzy Xor using other fuzzy connectives is not suitable for this purpose.

### 1. Introduction

The connective *exclusive or* plays an important role in computer programming especially in many encryption algorithms (see e.g. [9], [14], [15]), neural networks ([7], [13]), Quantum Computing ([10], [12]) or other fields ([4], [5], [16], [17]). The authors of [2] and [3] introduced an autonomous definition of the fuzzy Xor connective, which is independent of the other connectives. In the earlier literature we can find a fuzzy Xor operation as a composition of the fuzzy negation, and triangular conorms and norms (see [6], [8], [11], [13]).

In this paper we show what conditions must be met by the fuzzy Xor to be used for cryptographic purposes. We also provide new constructions of fuzzy Xor based on the composition of fuzzy implications and other fuzzy connectives.

### 2. Fuzzy connectives

In this article we use the notation  $\stackrel{(property)}{=}$ , which means that this equality holds due to the indicated property.

First, we recall definitions of some fuzzy connectives.

---

*Received:* 07.07.2025. *Accepted:* 26.10.2025.

(2020) Mathematics Subject Classification: 03B52.

*Key words and phrases:* fuzzy Xor, fuzzy implication, fuzzy negation, t-norm, t-conorm.

©2025 The Author(s).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License CC BY (<http://creativecommons.org/licenses/by/4.0/>).

DEFINITION 2.1. A *t-norm* is a function  $T: [0, 1]^2 \rightarrow [0, 1]$  satisfying

- (T1)  $T(x, y) = T(y, x)$ ,  $x, y \in [0, 1]$ ;
- (T2)  $T(x, T(y, z)) = T(T(x, y), z)$ ,  $x, y, z \in [0, 1]$ ;
- (T3)  $T(x, z) \leq T(y, z)$ ,  $x, y, z \in [0, 1]$ ,  $x \leq y$ ;
- (T4)  $T(x, 1) = x$ ,  $x \in [0, 1]$ .

DEFINITION 2.2. A *t-conorm* is a function  $S: [0, 1]^2 \rightarrow [0, 1]$  satisfying

- (S1)  $S(x, y) = S(y, x)$ ,  $x, y \in [0, 1]$ ;
- (S2)  $S(x, S(y, z)) = S(S(x, y), z)$ ,  $x, y, z \in [0, 1]$ ;
- (S3)  $S(x, z) \leq S(y, z)$ ,  $x, y, z \in [0, 1]$ ,  $x \leq y$ ;
- (S4)  $S(x, 0) = x$ ,  $x \in [0, 1]$ .

DEFINITION 2.3. A non-increasing function  $N: [0, 1] \rightarrow [0, 1]$  is called a *fuzzy negation* if  $N(0) = 1$ ,  $N(1) = 0$ . A fuzzy negation  $N$  is called

- *strict* if it is strictly decreasing and continuous;
- *strong* if it is an involution, i.e.,  $N(N(x)) = x$ ,  $x \in [0, 1]$ .

Given a fuzzy negation  $N$ , a value  $e \in (0, 1)$  such that  $N(e) = e$  is said to be an *equilibrium point* of  $N$ . If the equilibrium point of  $N$  exists (e.g. when  $N$  is continuous), then it is unique.

DEFINITION 2.4. A function  $I: [0, 1]^2 \rightarrow [0, 1]$  is called a *fuzzy implication* if it satisfies the following properties:

- (I1)  $I(x_2, y) \leq I(x_1, y)$ ,  $x_1, x_2, y \in [0, 1]$ ,  $x_1 \leq x_2$ ;
- (I2)  $I(x, y_1) \leq I(x, y_2)$ ,  $x, y_1, y_2 \in [0, 1]$ ,  $y_1 \leq y_2$ ;
- (I3)  $I(0, 0) = 1$ ;
- (I4)  $I(1, 1) = 1$ ;
- (I5)  $I(1, 0) = 0$ .

The *natural negation* of a fuzzy implication  $I$  is defined by  $N_I(x) := I(x, 0)$ ,  $x \in [0, 1]$ .

A fuzzy implication can have many properties (see [1]) but we will remind here only those that we will need, so for a fuzzy implication  $I$  and a fuzzy negation  $N$  we have the following properties:

- (R-CP(N))  $I(x, N(y)) = I(y, N(x))$ ,  $x, y \in [0, 1]$ ;
- (L-CP(N))  $I(N(x), y) = I(N(y), x)$ ,  $x, y \in [0, 1]$ ;
- (NP)  $I(1, x) = x$ ,  $x \in [0, 1]$ ;
- (IP)  $I(x, x) = 1$ ,  $x \in [0, 1]$ .

B. Bedregal, R.H.S. Reiser and G.P. Dimuro in their two works [2], [3] tried to introduce an intrinsic definition of the connective fuzzy exclusive or.

DEFINITION 2.5. A function  $E: [0, 1]^2 \rightarrow [0, 1]$  is called a *fuzzy Xor* if it satisfies the following properties:

- (E0)  $E(0, 0) = E(1, 1) = 0$ ,  $E(1, 0) = E(0, 1) = 1$ ;
- (E1)  $E(x, y) = E(y, x)$ ,  $x, y \in [0, 1]$ ;
- (E2a) functions  $E(0, \cdot), E(\cdot, 0)$  are non-decreasing;
- (E2b) functions  $E(1, \cdot), E(\cdot, 1)$  are non-increasing.

REMARK 2.6. Let  $E: [0, 1]^2 \rightarrow [0, 1]$  satisfy (E0) and (E2b). Then  $N_E: [0, 1] \rightarrow [0, 1]$  given by

$$N_E(x) := E(x, 1), \quad x \in [0, 1]$$

is a fuzzy negation. When  $E$  is a fuzzy Xor, then it is called the natural fuzzy negation of the fuzzy Xor  $E$ .

In [2], [3] we can find also the following properties connected with a fuzzy Xor function  $E$ :

- (E3)  $E(x, x) \neq 1$ ,  $x \in [0, 1]$ ;
- (E4)  $E(0, x) = x$ ,  $x \in [0, 1]$ ;
- (E5)  $E(E(x, y), z) = E(x, E(y, z))$ ,  $x, y, z \in [0, 1]$ ;
- (E6)  $E(x, x) = 0$ ,  $x \in [0, 1]$ ;
- (E7)  $E(E(x, y), y) = x$ ,  $x, y \in [0, 1]$ ;
- (E9)  $N_E$  is a strict fuzzy negation;
- (E10)  $N_E$  is a strong fuzzy negation;
- (E13)  $E(N_E(x), x) = 1$ ,  $x \in [0, 1]$ ;
- (E15)  $N_E(E(x, y)) = E(x, N_E(y))$ ,  $x, y \in [0, 1]$ .

In this work we will pay special attention to the property (E7), which is related to encryption. The authors of [3] showed that some of the above properties can be obtained from others (e.g. obvious implications (E6)  $\Rightarrow$  (E3), (E10)  $\Rightarrow$  (E9), (E5)  $\Rightarrow$  (E15)) but they missed that some of them are incompatible.

LEMMA 2.7. Let  $E: [0, 1]^2 \rightarrow [0, 1]$  satisfy (E13) and there exists an equilibrium point of  $N_E$ . Then (E3) does not hold.

PROOF. Let  $e \in (0, 1)$  be such that  $N_E(e) = e$ . We have

$$E(e, e) = E(N_E(e), e) = 1.$$

□

THEOREM 2.8. Let  $E: [0, 1]^2 \rightarrow [0, 1]$  satisfy (E7). Then, the following relations hold:

- (i) (E2b)  $\Rightarrow ((E10) \wedge \neg(E15) \wedge \neg(E5))$ ;
- (ii) (E1)  $\Rightarrow$  (E13);

- (iii)  $(E4) \Leftrightarrow (E6)$ ;
- (iv)  $(E2a) \Rightarrow E(x, 0) = x \text{ for } x \in [0, 1]$ ;
- (v)  $(E2b) \wedge (E13) \Rightarrow (\neg(E3) \wedge \neg(E4) \wedge \neg(E6))$ ;
- (vi)  $(E1) \wedge (E2b) \Rightarrow (\neg(E2a) \wedge \neg(E3) \wedge \neg(E4) \wedge \neg(E6))$ ;
- (vii)  $(E1) \wedge (E2a) \Rightarrow (\neg(E2b) \wedge (E3) \wedge (E4) \wedge (E6))$ ;
- (viii)  $((E2a) \wedge (E2b) \wedge (E7)) \Rightarrow (E0)$ .

PROOF. (i) We have

$$x \stackrel{(E7)}{=} E(E(x, 1), 1) = N_E^2(x), \quad x \in [0, 1].$$

Hence  $N_E^2(0) = 0$  and  $N_E^2(1) = 1$ , so from (E2b) we get  $N_E(0) = 1$  and  $N_E(1) = 0$ .

Suppose that (E15) holds. Since  $N_E$  is strong, there exists  $y_0 \in (0, 1)$  such that  $y_0 = N_E(y_0)$ . Then for  $x_0 := E(1, y_0)$  we have

$$\begin{aligned} 0 = N_E(1) &\stackrel{(E7)}{=} N_E(E(E(1, y_0), y_0)) = N_E(E(x_0, y_0)) \\ &\stackrel{(E15)}{=} E(x_0, N_E(y_0)) = E(x_0, y_0) = E(E(1, y_0), y_0) \stackrel{(E7)}{=} 1, \end{aligned}$$

which gives us a contradiction. Hence (E15) and (E5) do not hold.

(ii) We observe that

$$E(N_E(x), x) = E(E(x, 1), x) \stackrel{(E1)}{=} E(E(1, x), x) \stackrel{(E7)}{=} 1, \quad x \in [0, 1].$$

(iii) We have

$$E(x, x) \stackrel{(E4)}{=} E(E(0, x), x) \stackrel{(E7)}{=} 0, \quad x \in [0, 1]$$

and

$$E(0, x) \stackrel{(E6)}{=} E(E(x, x), x) \stackrel{(E7)}{=} x, \quad x \in [0, 1].$$

(iv) Let  $f(x) := E(x, 0)$ ,  $x \in [0, 1]$ . We notice that

$$f^2(x) = E(E(x, 0), 0) \stackrel{(E7)}{=} x, \quad x \in [0, 1],$$

so since  $f$  is non-decreasing,  $f(x) = x$ ,  $x \in [0, 1]$ .

(v) From (i) and Lemma 2.7 we know that (E3) does not hold, so (E6) does not hold. Hence from (iii) (E4) does not hold.

(vi) From (ii) and (v) we obtain that (E3), (E4), (E6) do not hold.

Suppose that (E2a) holds. We have  $E(0, x) \stackrel{(E1)}{=} E(x, 0) \stackrel{(iv)}{=} x$ ,  $x \in [0, 1]$ , so (E4) holds and we obtain the contradiction.

- (vii) We have  $E(0, x) \stackrel{(E1)}{=} E(x, 0) \stackrel{(iv)}{=} x$ ,  $x \in [0, 1]$ , so (E4) holds. From (iii) (E6) (and (E3)) holds. From (vi) (E2b) does not hold.
- (viii) We have

$$\begin{aligned}
 0 &\stackrel{(E7)}{=} E(E(0, 0), 0) \stackrel{(E2a)}{\geq} E(0, 0), \\
 1 &\stackrel{(E7)}{=} E(E(1, 0), 0) \stackrel{(E2a)}{\leq} E(1, 0), \\
 1 &\stackrel{(E7)}{=} E(E(1, 1), 1) \stackrel{(E2b)}{\leq} E(0, 1), \\
 0 &\stackrel{(E7)}{=} E(E(0, 1), 1) \stackrel{(E2b)}{\geq} E(1, 1). \quad \square
 \end{aligned}$$

REMARK 2.9. Let

(E1-0)  $E(x, 0) = E(0, x)$ ,  $x \in [0, 1]$ ;

(E1-1)  $E(x, 1) = E(1, x)$ ,  $x \in [0, 1]$ .

In the previous theorem instead of (E1) we can assume in (ii) that (E1-1) holds, in (vi) and (vii) that (E1-0), (E1-1) hold. Moreover, we have  $(E2a) \wedge (E4) \Rightarrow (E1-0)$ .

REMARK 2.10. There is no function that satisfies (E0), (E1), (E2a), (E2b) and (E7). Since conditions (E2a) and (E2b) are similar, then among functions satisfying (E7) it is better to look for these which do not satisfy (E1).

REMARK 2.11. Let  $E: [0, 1]^2 \rightarrow [0, 1]$  satisfy (E2a), (E2b) and (E7). Then we have the following possibilities:

- $E$  satisfies (E3), (E4), (E6) and does not satisfy (E13).
- $E$  satisfies (E13) and does not satisfy (E3), (E4), (E6).
- $E$  does not satisfy (E3), (E4), (E6), (E13).

All of the above possibilities can occur, as illustrated by the following examples.

EXAMPLE 2.12. Let  $N_1, N_2: [0, 1] \rightarrow [0, 1]$  be fuzzy negations,  $N_1$  be strong,  $E: [0, 1]^2 \rightarrow [0, 1]$  be a function given by

$$E(x, y) = \begin{cases} N_2(y) + (1 - N_2(y))N_1\left(\frac{x - N_2(y)}{1 - N_2(y)}\right), & x \geq N_2(y) \neq 1, \\ x, & x < N_2(y) \vee N_2(y) = 1. \end{cases}$$

We will show that  $E$  satisfies (E2a), (E2b), (E7) and does not satisfy (E3), (E4), (E6). Moreover (E13) holds iff  $N_1 = N_2$ . We have

$$E(x, 0) = x, \quad x \in [0, 1], \quad E(0, y) = \begin{cases} 1, & N_2(y) = 0, \\ 0, & N_2(y) \neq 0, \end{cases}$$

so (E2a) holds and (E4) does not hold,

$$\begin{aligned} N_E(x) &= E(x, 1) = N_1(x), & x \in [0, 1], \\ E(1, y) &= N_2(y), & y \in [0, 1], \end{aligned}$$

so (E2b) holds. Finally, for  $x, y \in [0, 1]$  we have

- if  $x \geq N_2(y) \neq 1$ , then  $E(x, y) \geq N_2(y)$  and

$$\begin{aligned} E(E(x, y), y) &= N_2(y) + (1 - N_2(y))N_1\left(\frac{E(x, y) - N_2(y)}{1 - N_2(y)}\right) \\ &= N_2(y) + (1 - N_2(y))N_1\left(\frac{N_2(y) + (1 - N_2(y))N_1\left(\frac{x - N_2(y)}{1 - N_2(y)}\right) - N_2(y)}{1 - N_2(y)}\right) \\ &= N_2(y) + (1 - N_2(y))N_1\left(N_1\left(\frac{x - N_2(y)}{1 - N_2(y)}\right)\right) \\ &= N_2(y) + (1 - N_2(y))\frac{x - N_2(y)}{1 - N_2(y)} = x, \end{aligned}$$

- if  $x < N_2(y) \vee N_2(y) = 1$ , then  $E(x, y) = x$  and

$$E(E(x, y), y) = E(x, y) = x,$$

so (E7) holds. From Theorem 2.8 (iii) (E6) and (E3) do not hold.

Now, we notice that

$$E(N_2(y), y) = 1,$$

so (E13)  $\Leftrightarrow N_1 = N_2$ .

EXAMPLE 2.13. Let  $N_1, N_2: [0, 1] \rightarrow [0, 1]$  be fuzzy negations,  $N_1$  be strong,  $N_2$  has not an equilibrium point. For  $y \in (0, 1)$  let  $A_y := (0, 1) \setminus \{y, N_2(y)\}$  and  $\varphi_y: A_y \rightarrow A_y$  be a bijective involution. Let further  $E: [0, 1]^2 \rightarrow [0, 1]$  be a function given by

$$E(x, y) = \begin{cases} \varphi_y(x), & y \in (0, 1), x \in A_y, \\ y, & x = 0, \\ N_2(y), & x = 1, \\ 0, & x = y \in (0, 1), \\ 1, & x = N_2(y) \in (0, 1), \\ x, & y = 0, x \in (0, 1), \\ N_1(x), & y = 1, x \in (0, 1). \end{cases}$$

We will show that  $E$  satisfies (E2a), (E2b), (E3), (E4), (E6), (E7) and does not satisfy (E13).

First, we observe that

$$\begin{aligned}
 E(E(x, y), y) &= \begin{cases} E(\varphi_y(x), y), & y \in (0, 1), x \in A_y, \\ E(y, y), & x = 0, \\ E(N_2(y), y), & x = 1, \\ E(0, y), & x = y \in (0, 1), \\ E(1, y), & x = N_2(y) \in (0, 1), \\ E(x, y), & y = 0, x \in (0, 1), \\ E(N_1(x), y), & y = 1, x \in (0, 1). \end{cases} \\
 &= \begin{cases} \varphi_y(\varphi_y(x)), & y \in (0, 1), x \in A_y, \\ 0, & x = 0, \\ 1, & x = 1, \\ y, & x = y \in (0, 1), \\ N_2(y), & x = N_2(y) \in (0, 1), \\ x, & y = 0, x \in (0, 1), \\ N_1(N_1(x)), & y = 1, x \in (0, 1). \end{cases} = x, \quad x, y \in [0, 1],
 \end{aligned}$$

so  $E$  satisfies (E7). We have also

$$\begin{aligned}
 E(x, 0) &= x, \quad x \in [0, 1], \\
 E(0, y) &= y, \quad y \in [0, 1], \\
 E(x, 1) &= N_1(x), \quad x \in [0, 1], \\
 E(1, y) &= N_2(y), \quad y \in [0, 1],
 \end{aligned}$$

whence (E2a), (E4), (E2b) hold. From Theorem 2.8 we obtain that (E6), (E3) hold and (E13) does not hold.

### 3. Fuzzy Xor generated from the other fuzzy connectives

In [2] and [3] we can find some classes of functions:

$$\begin{aligned}
 E_{S,T,N}(x, y) &= S\left(T(x, N(y)), T(N(x), y)\right), \quad x, y \in [0, 1]; \\
 E_{T,S,N}(x, y) &= T\left(S(x, y), S(N(x), N(y))\right), \quad x, y \in [0, 1]; \\
 E_{S,T}(x, y) &= S(x, y) - T(x, y), \quad x, y \in [0, 1];
 \end{aligned}$$

where  $T: [0, 1]^2 \rightarrow [0, 1]$  is a t-norm,  $S: [0, 1]^2 \rightarrow [0, 1]$  is a t-conorm, and  $N: [0, 1] \rightarrow [0, 1]$  is a fuzzy negation. Each of them is a fuzzy Xor satisfying (E4). We would like to present others classes which may not have this property.

### 3.1. Fuzzy Xor generated from fuzzy implications and negation

In a classical logic we can obtain a Xor in the following way

$$\alpha \oplus \beta = ((\alpha \Rightarrow \beta) \Rightarrow \neg(\beta \Rightarrow \alpha)).$$

Hence, we can define the following class of functions.

**DEFINITION 3.1.** Let  $I: [0, 1]^2 \rightarrow [0, 1]$  be a fuzzy implication,  $N: [0, 1] \rightarrow [0, 1]$  be a fuzzy negation. We define the function  $E_{I,N}: [0, 1]^2 \rightarrow [0, 1]$  by the formula

$$(1) \quad E_{I,N}(x, y) = I(I(x, y), N(I(y, x))), \quad x, y \in [0, 1].$$

We can generalize the above definition in the following way:

**DEFINITION 3.2.** Let  $I_1, I_2: [0, 1]^2 \rightarrow [0, 1]$  be fuzzy implications and  $N: [0, 1] \rightarrow [0, 1]$  be a fuzzy negation. We define the function  $E_{I_1, I_2, N}: [0, 1]^2 \rightarrow [0, 1]$  by the formula

$$(2) \quad E_{I_1, I_2, N}(x, y) = I_1(I_2(x, y), N(I_2(y, x))), \quad x, y \in [0, 1].$$

**REMARK 3.3.** Let  $S: [0, 1]^2 \rightarrow [0, 1]$  be a t-conorm,  $T: [0, 1]^2 \rightarrow [0, 1]$  be a t-norm,  $N: [0, 1] \rightarrow [0, 1]$  be a fuzzy negation. Let further  $\tilde{N}: [0, 1] \rightarrow [0, 1]$  be any strong negation. We define

$$\begin{aligned} I_1(x, y) &:= S(\tilde{N}(x), y), \quad x, y \in [0, 1], \\ I_2(x, y) &:= \tilde{N}(T(x, N(y))), \quad x, y \in [0, 1]. \end{aligned}$$

Then

$$\begin{aligned} E_{I_1, I_2, \tilde{N}}(x, y) &= I_1(I_2(x, y), \tilde{N}(I_2(y, x))) \\ &= S\left(\tilde{N}\left(\tilde{N}(T(x, N(y)))\right), \tilde{N}\left(\tilde{N}(T(y, N(x)))\right)\right) \\ &= S\left(T(x, N(y)), T(N(x), y)\right) = E_{S, T, N}(x, y), \quad x, y \in [0, 1]. \end{aligned}$$

**THEOREM 3.4.** Let  $I_1, I_2: [0, 1]^2 \rightarrow [0, 1]$  be fuzzy implications,  $N: [0, 1] \rightarrow [0, 1]$  be a fuzzy negation. Then for  $E_{I_1, I_2, N}$  given by (2) we have:

(i) (E0), (E2a), (E2b) hold and

$$\begin{aligned} (3) \quad & E_{I_1, I_2, N}(x, 0) = (N_{I_1} \circ N_{I_2})(x), \quad x \in [0, 1], \\ (4) \quad & E_{I_1, I_2, N}(0, x) = (g_{I_1} \circ N \circ N_{I_2})(x), \quad x \in [0, 1], \\ (5) \quad & E_{I_1, I_2, N}(x, 1) = (g_{I_1} \circ N \circ g_{I_2})(x), \quad x \in [0, 1], \\ (6) \quad & E_{I_1, I_2, N}(1, x) = (N_{I_1} \circ g_{I_2})(x), \quad x \in [0, 1], \end{aligned}$$

where  $g_{I_1}(x) := I_1(1, x)$ ,  $g_{I_2}(x) := I_2(1, x)$  for  $x \in [0, 1]$ ;



- (ii) if  $I_1$  satisfies (R-CP(N)), then (E1) holds, (E7) does not hold;
- (iii) (E4) holds iff  $(g_{I_1} \circ N \circ N_{I_2})(x) = x$  for  $x \in [0, 1]$ ;
- (iv) if  $I_1, I_2$  satisfy (R-CP(N)), then  $N_{E_{I_1, I_2, N}} \circ N = N_{I_1} \circ N_{I_2}$ ;
- (v) if  $I_1, I_2$  satisfy (R-CP(N)),  $N_{I_1}, N_{I_2}, N$  are strict, then (E9) holds;
- (vi) if  $I_2$  satisfies (IP), then (E6) holds;
- (vii) if  $I_1$  satisfies (R-CP(N)) and  $I_2$  satisfies (IP), then

$$E_{I_1, I_2, N}(x, y) = N_{I_1} \left( \min \left( I_2(x, y), I_2(y, x) \right) \right), \quad x, y \in [0, 1].$$

PROOF. (i) We observe that for  $x \in [0, 1]$  we have

$$\begin{aligned} E_{I_1, I_2, N}(x, 0) &= I_1 \left( I_2(x, 0), N(I_2(0, x)) \right) = I_1 \left( N_{I_2}(x), N(1) \right) \\ &= I_1(N_{I_2}(x), 0) = (N_{I_1} \circ N_{I_2})(x), \\ E_{I_1, I_2, N}(0, x) &= I_1 \left( I_2(0, x), N(I_2(x, 0)) \right) = I_1 \left( 1, N(N_{I_2}(x)) \right) \\ &= (g_{I_1} \circ N \circ N_{I_2})(x), \\ E_{I_1, I_2, N}(x, 1) &= I_1 \left( I_2(x, 1), N(I_2(1, x)) \right) = I_1 \left( 1, N(I_2(1, x)) \right) \\ &= (g_{I_1} \circ N \circ g_{I_2})(x), \\ E_{I_1, I_2, N}(1, x) &= I_1 \left( I_2(1, x), N(I_2(x, 1)) \right) = I_1 \left( I_2(1, x), N(1) \right) \\ &= I_1(I_2(1, x), 0) = (N_{I_1} \circ g_{I_2})(x), \end{aligned}$$

so we get (E0), (E2a), (E2b).

- (ii) Assume that  $I_1$  satisfies (R-CP(N)). Hence, using the above, we get

$$\begin{aligned} E_{I_1, I_2, N}(y, x) &= I_1 \left( I_2(y, x), N(I_2(x, y)) \right) \\ &\stackrel{(\text{R-CP(N)})}{=} I_1 \left( I_2(x, y), N(I_2(y, x)) \right) \\ &= E_{I_1, I_2, N}(x, y), \quad x, y \in [0, 1]. \end{aligned}$$

Using Theorem 2.8 we get that (E7) does not hold.

- (iii) It follows from i.
- (iv) Assume that  $I_2$  satisfies (R-CP(N)). Then we have

$$\begin{aligned} N_{E_{I_1, I_2, N}}(N(x)) &\stackrel{ii}{=} N_{I_1} \left( I_2(1, N(x)) \right) \stackrel{(\text{R-CP(N)})}{=} N_{I_1} \left( I_2(x, N(1)) \right) \\ &= N_{I_1} \left( I_2(x, 0) \right) = (N_{I_1} \circ N_{I_2})(x), \quad x \in [0, 1]. \end{aligned}$$

- (v) It follows from iv.

(vi) Assume that  $I_2$  satisfies (IP). Then

$$\begin{aligned} E_{I_1, I_2, N}(x, x) &= I_1(I_2(x, x), N(I_2(x, x))) \\ &\stackrel{(\text{IP})}{=} I_1(1, N(1)) = I_1(1, 0) = 0, \quad x \in [0, 1], \end{aligned}$$

so (E6) holds.

(vii) Let  $x, y \in [0, 1]$ . From ii we have that  $E_{I_1, I_2, N}$  is symmetric, so we can assume that  $y \leq x$  whence from (IP) for  $I_2$  we get  $I_2(y, x) = 1$  and

$$\begin{aligned} E_{I_1, I_2, N}(x, y) &= I_1(I_2(x, y), N(I_2(y, x))) \\ &= I_1(I_2(x, y), N(1)) = I_1(I_2(x, y), 0) = N_{I_1}(I_2(x, y)) \\ &= N_{I_1}(\min(I_2(x, y), I_2(y, x))). \end{aligned} \quad \square$$

LEMMA 3.5. *Let  $I: [0, 1]^2 \rightarrow [0, 1]$  be a fuzzy implication,  $N: [0, 1] \rightarrow [0, 1]$  be a fuzzy negation. Assume that  $E_{I, N}$  is given by (1). Then (E7) does not hold.*

PROOF. Suppose that (E7) holds. First, observe that  $N_I$  is strong. Let  $x \in [0, 1]$ . Using Theorem 2.8 (iv) we have

$$x = E_{I, N}(x, 0) \stackrel{(3)}{=} N_I(N_I(x)),$$

so  $N_I$  is strong.

Next, for  $x \in [0, 1]$  we notice that

$$N_{E_{I, N}}(x) = E_{I, N}(x, 1) \stackrel{(5)}{=} (g_I \circ N \circ g_I)(x).$$

Since  $g_I$  is non-decreasing,  $N_{E_{I, N}}$  is a strong fuzzy negation, then  $g_I$  is a bijection and  $N = g_I^{-1} \circ N_{E_{I, N}} \circ g_I^{-1}$  is strict.

The function  $g_I^{-1} \circ N_I \circ g_I \circ N \circ N_I$  is a strict fuzzy negation, so it has an equilibrium point  $e \in (0, 1)$ . Hence

$$(g_I \circ N \circ N_I)(e) = (N_I \circ g_I)(e).$$

Using the previous theorem we get

$$\begin{aligned} 1 &\stackrel{(\text{E7})}{=} E(E(1, e), e) \stackrel{(6)}{=} E((N_I \circ g_I)(e), e) \\ &= E((g_I \circ N \circ N_I)(e), e) \stackrel{(4)}{=} E(E(0, e), e) \stackrel{(\text{E7})}{=} 0, \end{aligned}$$

which gives us a contradiction.  $\square$

PROBLEM 3.6. Do exist fuzzy implications  $I_1, I_2$  and a negation  $N$  such that  $E_{I_1, I_2, N}$  satisfies (E7)?

### 3.2. Fuzzy Xor generated from t-norm, fuzzy implication and negation

In a classical logic we can obtain a Xor in the following way

$$\alpha \oplus \beta = ((\alpha \Rightarrow \neg\beta) \wedge (\neg\alpha \Rightarrow \beta)).$$

Hence, we can define the following class of functions.

**DEFINITION 3.7.** Let  $T: [0, 1]^2 \rightarrow [0, 1]$  be a t-norm,  $I: [0, 1]^2 \rightarrow [0, 1]$  be a fuzzy implication,  $N: [0, 1] \rightarrow [0, 1]$  be a fuzzy negation. We define the function  $E_{T,I,N}: [0, 1]^2 \rightarrow [0, 1]$  by the formula

$$(7) \quad E_{T,I,N}(x, y) = T\left(I(x, N(y)), I(N(x), y)\right), \quad x, y \in [0, 1].$$

**REMARK 3.8.** Let  $S: [0, 1]^2 \rightarrow [0, 1]$  be a t-conorm,  $T: [0, 1]^2 \rightarrow [0, 1]$  be a t-norm,  $N: [0, 1] \rightarrow [0, 1]$  be a strong fuzzy negation. We define

$$I(x, y) := S(N(x), y), \quad x, y \in [0, 1].$$

Then

$$\begin{aligned} E_{T,I,N}(x, y) &= T\left(I(x, N(y)), I(N(x), y)\right) \\ &= T\left(S(N(x), N(y)), S(N^2(x), y)\right) \\ &= T\left(S(N(x), N(y)), S(x, y)\right) = E_{T,S,N}(x, y), \quad x, y \in [0, 1]. \end{aligned}$$

**THEOREM 3.9.** Let  $T: [0, 1]^2 \rightarrow [0, 1]$  be a t-norm,  $I: [0, 1]^2 \rightarrow [0, 1]$  be a fuzzy implication,  $N: [0, 1] \rightarrow [0, 1]$  be a fuzzy negation. Then for  $E_{T,I,N}$  given by (7) we have:

(i) (E0), (E2a), (E2b) hold and

$$(8) \quad E_{T,I,N}(x, 0) = (N_I \circ N)(x), \quad x \in [0, 1],$$

$$(9) \quad E_{T,I,N}(0, x) = g_I(x), \quad x \in [0, 1],$$

$$(10) \quad E_{T,I,N}(x, 1) = N_I(x), \quad x \in [0, 1],$$

$$(11) \quad E_{T,I,N}(1, x) = (g_I \circ N)(x), \quad x \in [0, 1],$$

where  $g_I(x) := I(1, x)$  for  $x \in [0, 1]$ ;

- (ii) if  $I$  satisfies (R-CP(N)) and (L-CP(N)), then (E1) holds;
- (iii)  $I$  satisfies (NP) iff (E4) holds;
- (iv)  $N_I$  is strict (strong) iff (E9) ((E10)) holds;
- (v) if  $I$  satisfies (IP) and  $N$  has an equilibrium point, then (E3) does not hold;
- (vi) (E7) does not hold.

PROOF. (i) We observe that for  $x \in [0, 1]$  we have

$$\begin{aligned}
 E_{T,I,N}(x, 0) &= T\left(I(x, N(0)), I(N(x), 0)\right) = T\left(I(x, 1), (N_I \circ N)(x)\right) \\
 &= T\left(1, (N_I \circ N)(x)\right) = (N_I \circ N)(x), \\
 E_{T,I,N}(0, x) &= T\left(I(0, N(x)), I(N(0), x)\right) = T\left(1, I(1, x)\right) \\
 &= I(1, x) = g_I(x), \\
 E_{T,I,N}(x, 1) &= T\left(I(x, N(1)), I(N(x), 1)\right) \\
 &= T\left(I(x, 0), 1\right) = I(x, 0) = N_I(x), \\
 E_{T,I,N}(1, x) &= T\left(I(1, N(x)), I(N(1), x)\right) = T\left((g_I \circ N)(x), I(0, x)\right) \\
 &= T\left((g_I \circ N)(x), 1\right) = (g_I \circ N)(x),
 \end{aligned}$$

so we get (E0), (E2a), (E2b).

(ii) Assume that  $I$  satisfies (R-CP(N)) and (L-CP(N)). Hence

$$\begin{aligned}
 E_{T,I,N}(y, x) &= T\left(I(y, N(x)), I(N(y), x)\right) \\
 &\stackrel{\text{(R-CP(N))}}{=} T\left(I(x, N(y)), I(N(y), x)\right) \\
 &\stackrel{\text{(L-CP(N))}}{=} T\left(I(x, N(y)), I(N(x), y)\right) = E_{T,I,N}(x, y), \quad x, y \in [0, 1].
 \end{aligned}$$

(iii) It follows from (9).

(iv) It follows from (10).

(v) Assume that  $I$  satisfies (IP) and  $N$  has an equilibrium point  $e \in (0, 1)$ . Then

$$\begin{aligned}
 E_{T,I,N}(e, e) &= T\left(I(e, N(e)), I(N(e), e)\right) = T\left(I(e, e), I(e, e)\right) \\
 &\stackrel{\text{(IP)}}{=} T(1, 1) = 1, \quad x \in [0, 1].
 \end{aligned}$$

(vi) Suppose that (E7) holds. From Theorem 2.8  $N_{E_{T,I,N}}$  is strong, so from iv  $N_I$  is strong. From (8) we get

$$x \stackrel{\text{(E7)}}{=} E_{T,I,N}(E_{T,I,N}(x, 0), 0) \stackrel{\text{(8)}}{=} (N_I \circ N)^2(x), \quad x \in [0, 1].$$

Since  $N_I \circ N$  is non-decreasing  $(N_I \circ N)(x) = x$  for  $x \in [0, 1]$ , so  $N = N_I$ . Let  $e \in (0, 1)$  be an equilibrium point of  $N_I$ . Using i we have

$$\begin{aligned}
 1 &\stackrel{\text{(E7)}}{=} E_{T,I,N}(E_{T,I,N}(1, e), e) \stackrel{\text{(9)}}{=} E_{T,I,N}((g_I \circ N)(e), e) = E_{T,I,N}(g_I(e), e), \\
 0 &\stackrel{\text{(E7)}}{=} E_{T,I,N}(E_{T,I,N}(0, e), e) \stackrel{\text{(11)}}{=} E_{T,I,N}(g_I(e), e),
 \end{aligned}$$

which gives us a contradiction. □

#### 4. Examples

In [3, Table 2] we can find classification of fuzzy Xor connectives. Adding new classes of fuzzy Xor we can show that all these classes are different.

	$E_{I,N}$	$E_{T,I,N}$	$E_{S,T,N}$	$E_{T,S,N}$
$E_C(x, y)$	$(I_{RS}, N)$	-	-	-
$E_1(x, y)$	-	$(T, I_{LK}, N_C)$	$(S_{LK}, T_M, N_C)$	$(T, S_{LK}, N_C)$
$E_\perp(x, y)$	$(I_1, N)$	$(T, I_0, N)$	-	-
$E_2(x, y)$	-	-	$(S_D, T, N_{D2})$	$(T, S_D, N_{D2})$

where  $N$  is an arbitrary fuzzy negation,  $T$  is an arbitrary t-norm,

$$\begin{aligned}
E_C(x, y) &= \begin{cases} 0, & x = y, \\ 1, & x \neq y, \end{cases} & E_1(x, y) &= \begin{cases} x + y, & x + y \leq 1, \\ 2 - x - y, & x + y > 1, \end{cases} \\
E_\perp(x, y) &= \begin{cases} 0, & |x - y| = 1, \\ 1, & |x - y| \neq 1, \end{cases} & E_2(x, y) &= \begin{cases} 0, & x = y = 1, \\ \max(x, y), & x = 0 \vee y = 0, \\ 1, & \text{otherwise,} \end{cases} \\
I_{RS}(x, y) &= \begin{cases} 0, & x > y, \\ 1, & x \leq y, \end{cases} & I_{LK}(x, y) &= \begin{cases} 1 - x + y, & x > y, \\ 1, & x \leq y, \end{cases} \\
I_1(x, y) &= \begin{cases} 0, & x = 1 \wedge y = 0, \\ 1, & x < 1 \vee y > 0, \end{cases} & I_0(x, y) &= \begin{cases} 0, & x > 0 \wedge y < 1, \\ 1, & x = 0 \vee y = 1, \end{cases} \\
S_{LK}(x, y) &= \min(1, x + y), \quad x, y \in [0, 1], \\
S_D(x, y) &= \begin{cases} \max(x, y), & x = 0 \vee y = 0, \\ 1, & x > 0 \wedge y > 0, \end{cases} \\
T_M(x, y) &= \min(x, y), \quad x, y \in [0, 1], \\
N_C(x) &= 1 - x, \quad x \in [0, 1], \quad N_{D2}(x) = \begin{cases} 0, & x = 1, \\ 1, & x < 1. \end{cases}
\end{aligned}$$

It is easy to show the form of  $E$  if we know the generators, so we show only that the above fuzzy Xors do not belong to some classes. From [3, Table 2] we know that  $E_C$  and  $E_\perp$  do not belong to classes  $E_{S,T,N}$ ,  $E_{T,S,N}$ .

- Suppose that  $E_C = E_{T,I,N}$ . Using (8)–(11) we have

$$(N_I \circ N)(x) = g_I(x) = \begin{cases} 0, & x = 0, \\ 1, & x > 0, \end{cases} \quad N_I(x) = (g_I \circ N)(x) = \begin{cases} 0, & x = 1, \\ 1, & x < 1. \end{cases}$$

Hence  $I = I_1$ . Let  $x = y \in (0, 1)$ . Then  $I(x, N(y)) = 1 = I(N(x), y)$  and

$$0 = E_C(x, y) = E_{T,I,N}(x, y) = T(1, 1) = 1,$$

which give us a contradiction.

- Suppose that  $E_1 = E_{I,N}$ . Using (3)–(6) we have

$$\begin{aligned} N_I^2(x) &= (g_I \circ N \circ N_I)(x) = x, \quad x \in [0, 1], \\ (N_I \circ g_I)(x) &= (g_I \circ N \circ g_I)(x) = 1 - x, \quad x \in [0, 1]. \end{aligned}$$

Hence  $N_I = g_I \circ N$  is strong negation and

$$x = N_I^2(x) = (g_I \circ N \circ g_I \circ N)(x) = 1 - N(x), \quad x \in [0, 1],$$

so  $N = N_C$ .

Suppose that  $I(x_0, y_0) = 1$  for some  $x_0 > 0, y_0 < 1$ . Then

$$g_I(N_c(I(y, x))) = E_1(x, y) = \begin{cases} x + y, & x + y \leq 1, \\ 2 - x - y, & x + y > 1, \end{cases} \quad x < x_0, y > y_0.$$

The left hand side of the above identity is non-increasing in  $x$  and non-decreasing in  $y$  which is impossible, so  $I(x_0, y_0) = 1$  only if  $x_0 = 0$  or  $y_0 = 1$ .

Now, we observe that

$$1 = E_1(0.5, 0.5) = I(I(0.5, 0.5), 1 - I(0.5, 0.5)), \quad x \in [0, 1],$$

so  $I(0.5, 0.5) = 0$ , whence

$$0 < N_I(0.5) = I(0.5, 0) \leq I(0.5, 0.5) = 0,$$

which give us a contradiction.

- Suppose that  $E_2 = E_{I,N}$ . Using (3)–(6) we have

$$\begin{aligned} N_I^2(x) &= (g_I \circ N \circ N_I)(x) = x, \quad x \in [0, 1], \\ (N_I \circ g_I)(x) &= (g_I \circ N \circ g_I)(x) = N_{D2}(x), \quad x \in [0, 1]. \end{aligned}$$

Hence  $N_I$  is strong,  $g_I$  is surjective, so  $N_{D2} = N_I \circ g_I$  is surjective, which give us a contradiction.

Suppose that  $E_2 = E_{T,I,N}$ . Using (8)–(11) we have

$$\begin{aligned} (N_I \circ N)(x) &= g_I(x) = x, \quad x \in [0, 1], \\ N_I(x) &= (g_I \circ N)(x) = N_{D2}(x), \quad x \in [0, 1]. \end{aligned}$$

Hence  $N_I$  is surjective, so  $N_{D2} = N_I$  is surjective, which give us a contradiction.

## References

- [1] M. Baczyński and B. Jayaram, *Fuzzy Implications*, Studies in Fuzziness and Soft Computing, Vol. 231, Springer-Verlag, Berlin, 2008.
- [2] B.C. Bedregal, R.H.S. Reiser, and G.P. Dimuro, *Xor-implications and E-implications: classes of fuzzy implications based on fuzzy Xor*, Electron. Notes Theor. Comput. Sci. **247** (2009), 5–18.
- [3] B.C. Bedregal, R.H.S. Reiser, and G.P. Dimuro, *Revisiting Xor-implications: classes of fuzzy (co)implications based on f-Xor (f-XNor) connectives*, Internat. J. Uncertain. Fuzziness Knowledge-Based Systems **21** (2013), no. 6, 899–925.
- [4] G. Bongiovanni and P. Penna, *XOR-based schemes for fast parallel IP lookups*, Theory Comput. Syst. **38** (2005), no. 4, 481–501.
- [5] E.G. Coddling and G. Horlick, *Application of AND and exclusive-or (XOR) logic operations to the identification of elemental emission spectra measured using a photodiode array direct reading spectrometer*, Appl. Spectrosc. **27** (1973), no. 5, 366–370.
- [6] M.M. Gupta, A. Kandel, W. Bandler, and J.B. Kiszka (Eds.), *Approximate Reasoning in Expert Systems*, North-Holland Publishing Co., Amsterdam, 1985.
- [7] Y. Horikawa, *Landscapes of basins of local minima in the XOR problem*, in: *Proceedings of 1993 International Conference on Neural Networks (IJCNN-93-Nagoya, Japan)*, IEEE, 1993, pp. 1677–1680.
- [8] L. Kohout and E. Kim, *Characterization of interval fuzzy logic systems of connectives by group transformations*, Reliab. Comput. **10** (2004), no. 4, 299–334.
- [9] A.G. Konheim, *Computer Security and Cryptography*, John Wiley & Sons, Inc., Hoboken, 2007.
- [10] M. Maeda, M. Suenaga, and H. Miyajima, *Qubit neuron according to quantum circuit for XOR problem*, Appl. Math. Comput. **185** (2007), no. 2, 1015–1025.
- [11] C.F. Mela and D.R. Lehmann, *Using fuzzy set theoretic techniques to identify preference rules from interactions in the linear model: an empirical study*, Fuzzy Sets and Systems **71** (1995), no. 2, 165–181.
- [12] N.D. Mermin, *Quantum Computer Science. An Introduction*, Cambridge University Press, Cambridge, 2007.
- [13] W. Pedrycz and G. Succi, *fXOR fuzzy logic networks*, Soft Comput. **7** (2002), no. 2, 115–120.
- [14] C. Shannon, *Communication theory of secrecy systems*, Bell System Tech. J. **28** (1949), 656–715.
- [15] W. Stallings, *Cryptography and Network Security. Principles and Practice*, Prentice Hall, Upper Saddle River, 1999.
- [16] H. Vandierendonck and K. De Bosschere, *XOR-based hash functions*, IEEE Trans. Comput. **54** (2005), no. 7, 800–812.
- [17] H. Vandierendonck, P. Manet, and J.-D. Legat, *Application-specific reconfigurable XOR-indexing to eliminate cache conflict misses*, in: *Proceedings of the Design Automation & Test in Europe Conference*, IEEE, 2006, pp. 357–362.