

ON STABILITY OF THE CAUCHY FUNCTIONAL
EQUATION IN GROUPOIDS

IMKE TOBORG, PETER VOLKMANN

Abstract. We give some stability results for the functional equation $a(xy) = a(x) + a(y)$, where $a: G \rightarrow E$, G being a groupoid and E a Banach space.

1. Introduction

Let G be a groupoid, i.e., G is a set and for all $x, y \in G$ we have a product $xy \in G$. Furthermore, let E be a Banach space; by θ we denote its zero element.

We consider the Cauchy equation

$$(1) \quad a(xy) = a(x) + a(y), \quad x, y \in G,$$

for functions $a: G \rightarrow E$; its solutions are called *additive* functions.

A subset V of E is *ideally convex* (Evgenij Arkad'evič Lifšic [4]), if for every bounded sequence d_1, d_2, d_3, \dots in V and for every numerical sequence $\alpha_1, \alpha_2, \alpha_3, \dots \geq 0$ such that $\sum_{k=1}^{\infty} \alpha_k = 1$ we get $\sum_{k=1}^{\infty} \alpha_k d_k \in V$.

Let us mention that a convex subset of E is ideally convex, provided it is closed, open or finite dimensional (cf. also Jacek Tabor [8], where the relation between ideally convex sets and stability of the Cauchy equation has been examined; for this relation see also Volkman [11]). Thus closed and open balls

Received: 01.06.2016. Accepted: 20.09.2016. Published online: 18.12.2016.

(2010) Mathematics Subject Classification: 20N02, 39B82.

Key words and phrases: Cauchy equation, stability, groupoids, Tabor groupoids.

in E are ideally convex. We denote them by $S(p; \rho)$ and $S^0(p; \rho)$, respectively ($p \in E$ being the centre and $\rho \geq 0$ the radius).

Now we consider triplets (G, E, V) , where G, E, V essentially are as described before. More precisely, we introduce the following hypothesis:

(H) G is a groupoid, E a Banach space and V a bounded ideally convex subset of E .

DEFINITION 1. For a triplet (G, E, V) according to (H) we say it has *property* (U), if for every $f: G \rightarrow E$ satisfying

$$(2) \quad f(xy) - f(x) - f(y) \in V, \quad x, y \in G,$$

there is an additive $a: G \rightarrow E$ such that

$$(3) \quad a(x) - f(x) \in V, \quad x \in G.$$

REMARK 1. Concerning the special case $V = S(\theta; \varepsilon)$ (where $\varepsilon > 0$) we have:

Conditions (2), (3) can be written as

$$\|f(xy) - f(x) - f(y)\| \leq \varepsilon, \quad x, y \in G,$$

$$\|a(x) - f(x)\| \leq \varepsilon, \quad x, y \in G,$$

respectively, and (U) implies the Hyers–Ulam stability of the Cauchy equation (1) (in the sense of Zenon Moszner [5, Definition 1]; in fact, (U) is equivalent to the Hyers–Ulam stability of (1), which can be seen by using the Theorem 1 below). Finally property (U) for one $\varepsilon > 0$ implies already (U) for all $\varepsilon > 0$.

DEFINITION 2. For $x \in G$, G being a groupoid, and $k = 0, 1, 2, \dots$, the powers x^{2^k} are recursively defined by

$$x^{2^0} = x^1 = x, \quad x^{2^{k+1}} = x^{2^k} x^{2^k}.$$

The following result is a stability theorem for the functional equation $h(x^2) = 2h(x)$; see Volkmann [12] for the proof.

THEOREM 1. Consider (G, E, V) according to (H) and let $f: G \rightarrow E$ satisfy

$$(4) \quad f(x^2) - 2f(x) \in V, \quad x \in G.$$

Then there is exactly one $h: G \rightarrow E$ such that

$$(5) \quad h(x^2) = 2h(x), \quad h(x) - f(x) \in V, \quad x \in G,$$

namely

$$(6) \quad h(x) = \lim_{n \rightarrow \infty} \frac{1}{2^n} f(x^{2^n}), \quad x \in G.$$

In the next section we use this theorem for a characterization of property (U) and we give some applications. The third section will be devoted to direct products of groupoids and the last one to some concluding remarks.

2. A characterization of (U) and some consequences

The following result gives a characterization of property (U).

THEOREM 2. Consider (G, E, V) according to (H) and let $f: G \rightarrow E$ satisfy (2). Then the following assertions are equivalent:

- (A) There is an additive $a: G \rightarrow E$ satisfying (3).
- (B) The function $h: G \rightarrow E$ (given by (6)) is additive.

PROOF. (2) implies (4), and therefore we can apply Theorem 1: There is exactly one $h: G \rightarrow E$ satisfying (5), and this function is given by (6).

(A) \Rightarrow (B): If (A) holds, then the additive function a has all the properties of h , which are stated in (5). The uniqueness of h gives $h = a$, and this proves (B).

(B) \Rightarrow (A): If h is additive, then $a = h$ obviously leads to (A). □

REMARK 2. If the assertions (A), (B) of Theorem 2 are true, then $a = h$.

THEOREM 3. Consider (G, E, V) according to (H).

- I) If (U) holds for $(G, E, S(\theta; \varepsilon))$ ($\varepsilon > 0$), then (U) also holds for (G, E, V) .
- II) If $\text{Int}V \neq \emptyset$ and (U) holds for (G, E, V) , then (U) also holds for the triplet $(G, E, S(\theta; \varepsilon))$ ($\varepsilon > 0$).

PROOF. I) Let $f: G \rightarrow E$ satisfy (2). We choose $\varepsilon > 0$ such that $V \subseteq S(\theta; \varepsilon)$, and we get

$$(7) \quad f(xy) - f(x) - f(y) \in S(\theta; \varepsilon), \quad x, y \in G.$$

Since (U) holds for $(G, E, S(\theta; \varepsilon))$, we can apply Theorem 2 with V replaced by $S(\theta; \varepsilon)$ to get the additivity of $h: G \rightarrow E$ given by (6). This finishes the proof of (U) for (G, E, V) (because of (B) \Rightarrow (A) from Theorem 2).

II) We choose $p \in E$ and $\varepsilon > 0$ such that

$$S(p; \varepsilon) = p + S(\theta; \varepsilon) \subseteq V;$$

according to Remark 1 it is sufficient to keep the ε fixed and to show property (U) for $(G, E, S(\theta; \varepsilon))$. So let $f: G \rightarrow E$ satisfy (7). For

$$g(x) := f(x) - p, \quad x \in G,$$

we have

$$g(xy) - g(x) - g(y) = f(xy) - f(x) - f(y) + p \in S(p; \varepsilon) \subseteq V, \quad x, y \in G.$$

When using (U) for (G, E, V) , we get by Theorem 2 the additivity of

$$k(x) = \lim_{n \rightarrow \infty} \frac{1}{2^n} g(x^{2^n}), \quad x \in G.$$

For $h: G \rightarrow E$ given by (6) we now have

$$h(x) = \lim_{n \rightarrow \infty} \frac{1}{2^n} f(x^{2^n}) = \lim_{n \rightarrow \infty} \frac{1}{2^n} [g(x^{2^n}) + p] = k(x),$$

hence h is an additive function, and from Theorem 2 we get property (U) for $(G, E, S(\theta; \varepsilon))$. \square

The next definition is taken from Roman Badora, Barbara Przebieracz, Volkmann [1]; we adopt the notation $\mathbb{N} = \{0, 1, 2, \dots\}$, $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$.

DEFINITION 3. A groupoid G is called *Tabor groupoid*, if for $x, y \in G$ there exists $k \in \mathbb{N}^*$ such that

$$(8) \quad (xy)^{2^k} = x^{2^k} y^{2^k}.$$

Groups satisfying this condition had been considered by Józef Tabor [9]; we call them *Tabor groups*. The special case $k = 1$ in (8), i.e.,

$$(9) \quad (xy)^2 = x^2 y^2, \quad x, y \in G,$$

had been called square-symmetry by Zsolt Páles, Volkmann, R. Duncan Luce [6]. Of course, (9) holds in commutative semigroups.

The next theorem has two parts: Part I) is from Volkmann [12]; Part II) is similar to a result of Jürg Rätz [7].

THEOREM 4. *A triplet (G, E, V) according to (H) satisfies (U) in the following two cases:*

- I) G is a Tabor groupoid.
- II) For every $x \in G$ the set $\{x, x^2, x^4, x^8, \dots\}$ is finite.

PROOF. Let $f: G \rightarrow E$ satisfy (2). According to Theorem 2 it is sufficient to show the additivity of $h: G \rightarrow E$ (given by (6)). In Case I) this can be done by the procedure of Józef Tabor [9]. In Case II) we simply get $h(x) \equiv \theta$. \square

REMARK 3. For commutative semigroups G , Part I) goes back to Jacek Tabor [8].

REMARK 4. Condition II) is equivalent to the following: For every $x \in G$ there are $m, n \in \mathbb{N}$, $m \neq n$ such that $x^{2^m} = x^{2^n}$.

REMARK 5. In groupoids G , Rätz [7] uses the “left” powers, here for $x \in G$ written as $x^{(n)} := x(x^{(n-1)})$ ($n \in \mathbb{N} \setminus \{0, 1\}$), where $x^{(1)} := x$. By Rätz’ Theorem 2, a triplet (G, E, V) according to (H) also satisfies (U) in the following case:

- III) For every $x \in G$ the set $\{x^{(1)}, x^{(2)}, x^{(3)}, \dots\}$ is finite.

Let us give an example of a groupoid G , where I), II) hold but III) does not hold. We take $G = \mathbb{N}$, equipped with the product

$$x \circ y = \begin{cases} 0 & \text{if } x = y, \\ x + y + 1 & \text{if } x \neq y, \quad x, y \in \mathbb{N}. \end{cases}$$

I), II) are obviously satisfied, but

$$\begin{aligned} 1^{(2)} &= 1 \circ 1 = 0, & 1^{(3)} &= 1 \circ 0 = 2, & 1^{(4)} &= 1 \circ 2 = 4, \\ 1^{(n)} &= 2(n - 2), & n &\geq 2. \end{aligned}$$

$\{1^{(1)}, 1^{(2)}, 1^{(3)}, \dots\}$ is an infinite set, hence III) does not hold.

3. Direct products of groupoids

Let G, H be groupoids. By the direct product of them we understand (as usual) $G \times H$ equipped with the coordinate-wise defined product, i.e.,

$$(x, y)(\bar{x}, \bar{y}) = (x\bar{x}, y\bar{y}), \quad x, \bar{x} \in G; y, \bar{y} \in H.$$

A basic question is the following: Let furthermore V be an ideally convex set in a Banach space E , and suppose $(G, E, V), (H, E, V)$ to have property (U). Under which conditions is it true that $(G \times H, E, V)$ also has property (U)?

Concerning this question, Badora, Przebieracz, Volkmann [2] observed that $G \times H$ is a Tabor groupoid, provided G, H have this property and in at least one of them the square-symmetry (9) holds. This fact also follows from Theorem 5 below, which gives a necessary and sufficient condition for $G \times H$ to be a Tabor groupoid.

DEFINITION 4. For groupoids G and $x, y \in G$ we set

$$T_G(x, y) = \{k | k \in \mathbb{N}^*, (xy)^{2^k} = x^{2^k} y^{2^k}\}.$$

REMARK 6. A groupoid G is a Tabor groupoid if and only if

$$T_G(x, y) \neq \emptyset, \quad x, y \in G,$$

and in square-symmetric groupoids G we have

$$T_G = \mathbb{N}^*, \quad x, y \in G.$$

THEOREM 5. *Let G, H be groupoids. Then $G \times H$ is a Tabor groupoid if and only if $T_G(x, y) \cap T_H(a, b) \neq \emptyset$ ($x, y \in G; a, b \in H$). In particular G and H are Tabor groupoids in this case.*

PROOF. Consider $x, y \in G$ and $a, b \in H$. The theorem follows from the formula

$$T_{G \times H}((x, a), (y, b)) = T_G(x, y) \cap T_H(a, b),$$

which is easily shown: For $k \in T_{G \times H}((x, a), (y, b))$ we have

$$\begin{aligned} ((xy)^{2^k}, (ab)^{2^k}) &= (xy, ab)^{2^k} = ((x, a)(y, b))^{2^k} \\ &= (x, a)^{2^k} (y, b)^{2^k} = (x^{2^k}, a^{2^k})(y^{2^k}, b^{2^k}) = (x^{2^k} y^{2^k}, a^{2^k} b^{2^k}), \end{aligned}$$

hence $k \in T_G(x, y) \cap T_H(a, b)$. On the other hand, for $k \in T_G(x, y) \cap T_H(a, b)$, we get

$$\begin{aligned} ((x, a)(y, b))^{2^k} &= (xy, ab)^{2^k} = ((xy)^{2^k}, (ab)^{2^k}) = \\ &= (x^{2^k}y^{2^k}, a^{2^k}b^{2^k}) = (x^{2^k}, a^{2^k})(y^{2^k}, b^{2^k}) = (x, a)^{2^k}(y, b)^{2^k}, \end{aligned}$$

hence $k \in T_{G \times H}((x, a), (y, b))$. □

THEOREM 6. *Let (G, E, V) satisfy (U). Let Σ be a groupoid with an element $\sigma = \sigma^2 \in \Sigma$ such that for every $\xi \in \Sigma$ there exists $m \in \mathbb{N}$ yielding $\xi^{2^m} = \sigma$. Then $(G \times \Sigma, E, V)$ also has property (U).*

PROOF. Let $f: G \times \Sigma \rightarrow E$ satisfy

$$(10) \quad f(xy, \xi\eta) - f(x, \xi) - f(y, \eta) \in V, \quad x, y \in G; \xi, \eta \in \Sigma.$$

Then $y = x, \eta = \xi$ leads to

$$f(x^2, \xi^2) - 2f(x, \xi) \in V, \quad x \in G, \xi \in \Sigma,$$

and Theorem 1 applied to $(G \times \Sigma, E, V)$ shows the existence of exactly one function $h: G \times \Sigma \rightarrow E$ such that

$$(11) \quad h(x^2, \xi^2) = 2h(x, \xi), \quad h(x, \xi) - f(x, \xi) \in V, \quad x \in G, \xi \in \Sigma.$$

This function is given by

$$h(x, \xi) = \lim_{n \rightarrow \infty} \frac{1}{2^n} f(x^{2^n}, \xi^{2^n}), \quad x \in G, \xi \in \Sigma.$$

The choice $\xi = \eta = \sigma$ in (10) leads to

$$f(xy, \sigma) - f(x, \sigma) - f(y, \sigma) \in V, \quad x, y \in G,$$

and since (G, E, V) has the property (U), we get an additive $a: G \rightarrow E$ such that

$$a(x) - f(x, \sigma) \in V, \quad x \in G.$$

By Theorem 2 and Remark 2 we now have

$$(12) \quad a(x) = \lim_{n \rightarrow \infty} \frac{1}{2^n} f(x^{2^n}, \sigma) = h(x, \sigma), \quad x \in G.$$

For $\xi \in \Sigma$ let $m \in \mathbb{N}$ be such that $\xi^{2^m} = \sigma$. Then we get by (11), (12) for $x \in G$:

$$\begin{aligned} h(x, \xi) &= \frac{1}{2}h(x^2, \xi^2) = \dots \\ &= \frac{1}{2^m}h(x^{2^m}, \sigma) = \frac{1}{2^m}a(x^{2^m}) = a(x). \end{aligned}$$

So we have

$$h(x, \xi) = a(x), \quad x \in G, \xi \in \Sigma,$$

and therefore the function $h: G \times \Sigma \rightarrow E$ occurring in (11) is additive. Indeed, for $x, y \in G$ and $\xi, \eta \in \Sigma$ we get

$$\begin{aligned} h((x, \xi)(y, \eta)) &= h(xy, \xi\eta) = a(xy) \\ &= a(x) + a(y) = h(x, \xi) + h(y, \eta). \end{aligned}$$

This finishes the proof of (U) for $(G \times \Sigma, E, V)$. □

REMARK 7. When taking $G = \{0\}$ (a singleton) and observing $\Sigma \cong \{0\} \times \Sigma$, we see that (Σ, E, V) has property (U). In fact, Σ is a Tabor groupoid (a proof is easy), which in the semigroup-case already is known from Badora, Przebieracz, Volkmann [2, Theorem 3, Case I)].

The next theorem is trivial (we omit the proof), but it may be useful in applications.

THEOREM 7. *Let (G_1, E_1, V_1) , (G_2, E_2, V_2) have property (U) and let the function $f: G_1 \times G_2 \rightarrow E_1 \times E_2$ be given by*

$$f(x_1, x_2) = (f_1(x_1), f_2(x_2)), \quad (x_1, x_2) \in G_1 \times G_2,$$

where $f_j: G_j \rightarrow E_j$ ($j = 1, 2$). Suppose

$$f(xy) - f(x) - f(y) \in V_1 \times V_2, \quad x, y \in G_1 \times G_2.$$

Then there exists an additive $a: G_1 \times G_2 \rightarrow E_1 \times E_2$ such that

$$a(x) - f(x) \in V_1 \times V_2, \quad x \in G_1 \times G_2.$$

REMARK 8. If $E_1 \times E_2$ is normed by $\|(x_1, x_2)\| = \max\{\|x_1\|, \|x_2\|\}$ and V_1, V_2 are closed (or open) ε -balls centered at θ , then $V_1 \times V_2$ also is a closed (or open) ε -ball centered at θ , i.e.,

$$S_{E_1}(\theta; \varepsilon) \times S_{E_2}(\theta; \varepsilon) = S_{E_1 \times E_2}(\theta; \varepsilon),$$

$$S_{E_1}^0(\theta; \varepsilon) \times S_{E_2}^0(\theta; \varepsilon) = S_{E_1 \times E_2}^0(\theta; \varepsilon).$$

Of course, this remark concerns in particular the Hyers–Ulam stability mentioned in Remark 1.

Let us conclude this section by recalling some known results for groups.

THEOREM 8. *Let G be a group.*

- I) *If every element of G has odd order, then G is a Tabor group.*
- II) *If every element of G has an order 2^n (where $n \in \mathbb{N}$), then G is a Tabor group.*
- III) *If $G \cong G_1 \times G_2$ with groups G_1, G_2 as in I), II) (respectively), then G is a Tabor group.*
- IV) *Any finite Tabor group G has the form given in III).*

REMARK 9. I), II) follow from Badora, Przebieracz, Volkmann [2], concerning II) cf. Remark 7; III), IV) are from Toborg [10].

4. Final remarks

Let $F(a, b)$ be the free group with two generators and let \mathbb{R} denote the space of the reals. Gian Luigi Forti [3] has shown that the triplet $(F(a, b), \mathbb{R}, [-1, 1])$ does not have property (U). Thus $F(a, b)$ is not a Tabor group. Now the question is of interest, whether there exist torsion free non commutative Tabor groups.

Finally let us mention that all groupoids with two elements are Tabor groupoids; this can be easily checked. On the other hand, there is a groupoid $G = \{a, b, c\}$ which is not a Tabor groupoid: It is sufficient to require $a^2 = a, ab = c, b^2 = c^2 = b \neq c$. Indeed, assume for some $k \in \mathbb{N}^*$ that

$$(ab)^{2^k} = a^{2^k} b^{2^k}.$$

We get $c^{2^k} = ab$, hence $b = c$, which is a contradiction.

Acknowledgement. We cordially thank Mrs. Marion Ewald for her typesetting of this paper.

References

- [1] Badora R., Przebieracz B., Volkmann P., *Stability of the Pezider functional equation*, Ann. Math. Sil. **24** (2010), 7–13.
- [2] Badora R., Przebieracz B., Volkmann P., *On Tabor groupoids and stability of some functional equations*, Aequationes Math. **87** (2014), 165–171.
- [3] Forti G.L., *Remark 11* (at the 22nd International Symposium on Functional Equations, Oberwolfach 1984), Aequationes Math. **29** (1985), 90–91.
- [4] Lifšic E.A., *Ideal'no vypuklye množestva*, Funkcional'. Analiz Priložen. **4** (1970), no. 4, 76–77.
- [5] Moszner Z., *On the stability of functional equations*, Aequationes Math. **77** (2009), 33–88.
- [6] Páles Z., Volkmann P., Luce R.D., *Hyers–Ulam stability of functional equations with a square-symmetric operation*, Proc. Nat. Acad. Sci. U.S.A. **95** (1998), 12772–12775.
- [7] Rätz J., *On approximately additive mappings*, in: *General inequalities 2*, International Series of Numerical Mathematics 47, Birkhäuser, Basel, 1980, pp. 233–251.
- [8] Tabor Jacek, *Ideally convex sets and Hyers theorem*, Funkcial. Ekvac. **43** (2000), 121–125.
- [9] Tabor Józef, *Remark 18* (at the 22nd International Symposium on Functional Equations, Oberwolfach 1984), Aequationes Math. **29** (1985), 96.
- [10] Toborg I., *Tabor groups with finiteness conditions*, Aequationes Math. **90** (2016), 699–704.
- [11] Volkmann P., *Zur Rolle der ideal konvexen Mengen bei der Stabilität der Cauchyschen Funktionalgleichung*, Sem. LV, no. 6 (1999), 6 pp., <http://www.math.us.edu.pl/smdk>
- [12] Volkmann P., *O stabilności równań funkcyjnych o jednej zmiennej*, Sem. LV, no. 11 (2001), 6 pp., Errata ibid. no. 11bis (2003), 1 p., <http://www.math.us.edu.pl/smdk>

INSTITUT FÜR MATHEMATIK
NATURWISSENSCHAFTLICHE FAKULTÄT II
MARTIN-LUTHER-UNIVERSITÄT
HALLE-WITTENBERG
06099 HALLE (SAALE)
GERMANY
e-mail: imke.toborg@mathematik.uni-halle.de

INSTITUT FÜR ANALYSIS
KIT
76128 KARLSRUHE
GERMANY