

Annales Mathematicae Silesianae

26 (2012)

WYDAWNICTWO UNIWERSYTETU ŚLĄSKIEGO
KATOWICE 2013



ORTHOGONALLY PEXIDER FUNCTIONS MODULO A DISCRETE SUBGROUP

WIRGINIA WYROBEK-KOCHANEK

Abstract. Under appropriate conditions on abelian topological groups G and H , an orthogonality $\perp \subset G^2$ and a σ -algebra \mathfrak{M} of subsets of G we prove that if at least one of the functions $f, g, h: G \rightarrow H$ satisfying

$$f(x+y) - g(x) - h(y) \in K \quad \text{for } x, y \in G \text{ such that } x \perp y,$$

where K is a discrete subgroup of H , is continuous at a point or \mathfrak{M} -measurable, then there exist: a continuous additive function $A: G \rightarrow H$, a continuous biadditive and symmetric function $B: G \times G \rightarrow H$ and constants $a, b \in H$ such that

$$\begin{cases} f(x) - B(x, x) - A(x) - a \in K, \\ g(x) - B(x, x) - A(x) - b \in K, \\ h(x) - B(x, x) - A(x) - a + b \in K \end{cases}$$

for $x \in G$ and

$$B(x, y) = 0 \quad \text{for } x, y \in G \text{ such that } x \perp y.$$

Let G and H be groups and $\perp \subset G^2$ an orthogonality. We say that a function $f: G \rightarrow H$ is orthogonally additive, if

$$f(x+y) = f(x) + f(y) \quad \text{for } x, y \in G \text{ such that } x \perp y.$$

Received: 14.02.2012. *Revised:* 15.06.2012.

(2010) Mathematics Subject Classification: 39B55, 39B52.

Key words and phrases: additive functions, biadditive functions, Pexider difference, quadratic functions.

In the paper [3] J. Brzdęk considers the Rätz orthogonality (cf.[5]) and, under some assumptions, gives a description of orthogonally additive functions modulo a discrete subgroup, i.e. functions $f: G \rightarrow H$ such that

$$f(x+y) - f(x) - f(y) \in K \quad \text{for } x, y \in G \text{ such that } x \perp y,$$

where K is a discrete subgroup of H . In the papers [7] and [4] authors prove similar theorems (for continuous or measurable functions), but for the orthogonality defined by K. Baron and P. Volkman in [2], which includes the Rätz orthogonality.

Now we would like to obtain some similar results for the Pexider difference instead of the Cauchy difference, i.e. we assume that functions $f, g, h: G \rightarrow H$ are orthogonally Pexider modulo a discrete subgroup, which means that they satisfy

$$f(x+y) - g(x) - h(x) \in K \quad \text{for } x, y \in G \text{ such that } x \perp y,$$

where K is a discrete subgroup of H . We start with the following result.

LEMMA. *Let G be a groupoid with a neutral element, H an abelian group, K a subgroup of H . Let $\Delta \subset G \times G$ be a set with*

$$(1) \quad (0, x), (x, 0) \in \Delta \quad \text{for all } x \in G.$$

If functions $f, g, h: G \rightarrow H$ satisfy

$$(2) \quad f(x+y) - g(x) - h(y) \in K \quad \text{for } (x, y) \in \Delta,$$

then the following are true:

(a) *There are functions $k_1, l_1: G \rightarrow K$, $\varphi_1: G \rightarrow H$ and constants $a, b \in H$ such that*

$$\varphi_1(x+y) - \varphi_1(x) - \varphi_1(y) \in K \quad \text{for } (x, y) \in \Delta$$

and

$$(3) \quad \begin{cases} f(x) = \varphi_1(x) + a, \\ g(x) = \varphi_1(x) + k_1(x) + b, \\ h(x) = \varphi_1(x) - k_1(x) + l_1(x) + a - b \end{cases}$$

for all $x \in G$.

- (b) *There are functions $k_2, l_2: G \rightarrow K$, $\varphi_2: G \rightarrow H$ and constants $a, b \in H$ such that*

$$\varphi_2(x+y) - \varphi_2(x) - \varphi_2(y) \in K \quad \text{for } (x, y) \in \Delta$$

and

$$\begin{cases} f(x) = \varphi_2(x) + k_2(x) + a, \\ g(x) = \varphi_2(x) + b, \\ h(x) = \varphi_2(x) + l_2(x) + a - b \end{cases}$$

for all $x \in G$.

- (c) *There are functions $k_3, l_3: G \rightarrow K$, $\varphi_3: G \rightarrow H$ and constants $a, b \in H$ such that*

$$\varphi_3(x+y) - \varphi_3(x) - \varphi_3(y) \in K \quad \text{for } (x, y) \in \Delta$$

and

$$\begin{cases} f(x) = \varphi_3(x) + k_3(x) + a, \\ g(x) = \varphi_3(x) + l_3(x) + b, \\ h(x) = \varphi_3(x) + a - b \end{cases}$$

for all $x \in G$.

Moreover, each of assertions (a), (b), (c) gives a complete description of solutions of (2), that is, every triple (f, g, h) , being of one of the forms described above, is a solution of (2).

PROOF. Setting $y = 0$ in (2), by (1) we get

$$(4) \quad \mu(x) := f(x) - g(x) - h(0) \in K \quad \text{for } x \in G$$

and setting $x = 0$ we have

$$(5) \quad \nu(y) := f(y) - g(0) - h(y) \in K \quad \text{for } y \in G.$$

In particular,

$$(6) \quad f(0) - g(0) - h(0) \in K.$$

Denote $a = f(0)$, $b = g(0)$ and define $\varphi_i, k_i, l_i: G \rightarrow H$ for $i = 1, 2, 3$ by

$$\begin{aligned}\varphi_1 &= f - a, & k_1 &= g - \varphi_1 - b, & l_1 &= h + k_1 - \varphi_1 - a + b, \\ \varphi_2 &= g - b, & k_2 &= f - \varphi_2 - a, & l_2 &= h - \varphi_2 - a + b, \\ \varphi_3 &= h - a + b, & k_3 &= f - \varphi_3 - a, & l_3 &= g - \varphi_3 - b.\end{aligned}$$

Using (4), (5), (2) and (6) for every $(x, y) \in \Delta$ we get

$$\begin{aligned}\varphi_1(x+y) - \varphi_1(x) - \varphi_1(y) &= f(x+y) - a - f(x) + a - f(y) + a \\ &= f(x+y) - \mu(x) - g(x) - h(0) - \nu(y) - g(0) - h(y) + a \in K; \\ \varphi_2(x+y) - \varphi_2(x) - \varphi_2(y) &= g(x+y) - b - g(x) + b - g(y) + b \\ &= f(x+y) - \mu(x+y) - h(0) - g(x) + \mu(y) - f(y) + h(0) + b \\ &= f(x+y) - \mu(x+y) - g(x) + \mu(y) - \nu(y) - g(0) - h(y) + b \in K; \\ \varphi_3(x+y) - \varphi_3(x) - \varphi_3(y) &= h(x+y) - a + b - h(x) + a - b - h(y) + a - b \\ &= f(x+y) - g(0) - \nu(x+y) + \nu(x) - f(x) + g(0) - h(y) + a - b \\ &= f(x+y) - \nu(x+y) + \nu(x) - \mu(x) - g(x) - h(0) - h(y) + a - b, \\ &\in K.\end{aligned}$$

We also have

$$\begin{aligned}k_1(x) &= g(x) - f(x) + a - b = -\mu(x) - h(0) + a - b \in K, \\ k_2(x) &= f(x) - g(x) + b - a = \mu(x) + h(0) + b - a \in K, \\ k_3(x) &= f(x) - h(x) + a - b - a = \nu(x) + g(0) - b \in K, \\ l_1(x) &= h(x) + k_1(x) - f(x) + a - a + b = -\nu(x) - g(0) + k_1(x) + b \in K, \\ l_2(x) &= h(x) + k_2(x) - f(x) + a - a + b = -\nu(x) - g(0) + k_2(x) + b \in K, \\ l_3(x) &= g(x) + k_3(x) - f(x) + a - b = -\mu(x) - h(0) + k_3(x) + a - b \in K\end{aligned}$$

for $x \in G$. □

The part (b) of this lemma in the case when $\Delta = G^2$ was also obtained by K. Baron and PL. Kannappan in [1], even under some weaker assumptions. Some variations of (2) for functions with values in groupoids were studied by J. Sikorska in [6].

We work with the orthogonality proposed by K. Baron and P. Volkmann in [2], assuming additionally that the last condition in the following definition holds:

Let G be a group such that the mapping

$$(7) \quad x \mapsto 2x, \quad x \in G,$$

is a bijection onto the group G . A relation $\perp \subset G^2$ is called *orthogonality* if it satisfies the following three conditions:

- (i) $0 \perp 0$; and from $x \perp y$ the relations $-x \perp -y$, $\frac{x}{2} \perp \frac{y}{2}$ follow.
- (ii) If an orthogonally additive function from G to an abelian group is odd, then it is additive; if it is even, then it is quadratic.
- (iii) $x \perp 0$ and $0 \perp x$ for every $x \in G$.

For a subset U of a given group and for $n \in \mathbb{N}$ the symbol nU denotes the set $\{nx : x \in U\}$.

THEOREM. *Assume G is an abelian topological group such that the mapping (7) is a homeomorphism and every neighbourhood of zero in G contains a neighbourhood U of zero such that*

$$(8) \quad U \subset 2U \quad \text{and} \quad G = \bigcup \{2^n U : n \in \mathbb{N}\}.$$

Let $\perp \subset G^2$ be an orthogonality, H an abelian topological group and K a discrete subgroup of H . Assume that functions $f, g, h: G \rightarrow H$ satisfy

$$(9) \quad f(x+y) - g(x) - h(y) \in K \quad \text{for } x, y \in G \text{ such that } x \perp y.$$

(i) *If at least one of the functions f, g, h is continuous at a point, then there exist: a continuous additive function $A: G \rightarrow H$, a continuous biadditive and symmetric function $B: G \times G \rightarrow H$ and constants $a, b \in H$ such that*

$$(10) \quad \begin{cases} f(x) - B(x, x) - A(x) - a \in K, \\ g(x) - B(x, x) - A(x) - b \in K, \\ h(x) - B(x, x) - A(x) - a + b \in K \end{cases}$$

for $x \in G$ and

$$(11) \quad B(x, y) = 0 \quad \text{for } x, y \in G \text{ such that } x \perp y.$$

(ii) *Let \mathfrak{M} be a σ -algebra of subsets of G such that*

$$(12) \quad x \pm 2A \in \mathfrak{M} \quad \text{for all } x \in G \text{ and } A \in \mathfrak{M}$$

and there is a proper σ -ideal \mathfrak{J} of subsets of G with

$$(13) \quad 0 \in \text{Int}(A - A) \quad \text{for } A \in \mathfrak{M} \setminus \mathfrak{J}.$$

Assume moreover that H is separable metric and the following condition (G) is fulfilled:

(G) either G is a first countable Baire group, or G is metric separable, or G is metric and \mathfrak{M} contains all Borel subsets of G .

If at least one of the functions f, g, h is \mathfrak{M} -measurable, then there exist: a continuous additive function $A: G \rightarrow H$, a continuous biadditive and symmetric function $B: G \times G \rightarrow H$ and constants $a, b \in H$ such that (10) and (11) hold.

Moreover, each of assertions (i), (ii) gives a complete description of solutions of (9).

PROOF. (i): Case 1. Assume that f is continuous at a point. Let $k_1, l_1: G \rightarrow K$, $\varphi_1: G \rightarrow H$ be as in Lemma (a). Then the function φ_1 is continuous at a point. According to Theorem 1 from [7] we get a continuous additive function $A: G \rightarrow H$ and a continuous biadditive and symmetric function $B: G \times G \rightarrow H$ such that

$$\varphi_1(x) - B(x, x) - A(x) \in K \quad \text{for } x \in G$$

and (11) hold. Then, according to (3),

$$\begin{aligned} f(x) - B(x, x) - A(x) - a &= \varphi_1(x) + a - B(x, x) - A(x) - a \in K, \\ g(x) - B(x, x) - A(x) - b &= \varphi_1(x) + k_1(x) + b - B(x, x) - A(x) - b \in K, \\ h(x) - B(x, x) - A(x) - a + b &= \varphi_1(x) - k_1(x) + l_1(x) + a - b \\ &\quad - B(x, x) - A(x) - a + b \in K \end{aligned}$$

for all $x \in G$.

Case 2. If the function g is continuous at a point then instead of Lemma (a) we use Lemma (b).

Case 3. If the function h is continuous at a point then we use Lemma (c).

(ii): If one of the functions f, g, h is \mathfrak{M} -measurable then we use Theorem 1 from [4] instead of Theorem 1 from [7]. \square

For $\perp = G^2$ some special cases were obtained in [1] (cf. Corollaries 6 and 7 there).

If in the Theorem G is Baire and we consider the Baire measurability, then we do not need to assume the first countability of G in order to get the factorization with a separately continuous biadditive term only (cf. Corollary 2 in [4]).

COROLLARY 1. Assume G is an abelian topological group such that the mapping (7) is a homeomorphism and every neighbourhood of zero in G contains a neighbourhood U of zero such that (8) holds. Let $\perp \subset G^2$ be an

orthogonality, H an abelian separable metric group, K a discrete subgroup of H and functions $f, g, h: G \rightarrow H$ satisfy (9). If G is Baire and at least one of the functions f, g, h is Baire measurable, then there exist: a continuous additive function $A: G \rightarrow H$, a function $B: G \times G \rightarrow H$ biadditive, symmetric and continuous in each variable, and constants $a, b \in H$ such that (10) and (11) hold.

If we take $\perp = G^2$, then our Theorem gives us Corollary 2 below. It also leads to another conclusions in the case when we consider Baire or Christensen measurability.

COROLLARY 2. *Assume G is an abelian topological group such that the mapping (7) is a homeomorphism and every neighbourhood of zero in G contains a neighbourhood U of zero such that (8) holds. Let H be an abelian separable metric group, K a discrete subgroup of H , \mathfrak{M} a σ -algebra of subsets of G satisfying (12) and such that there is a proper σ -ideal \mathfrak{J} of subsets of G with property (13). If functions $f, g, h: G \rightarrow H$ satisfy*

$$f(x + y) - g(x) - h(y) \in K \quad \text{for } x, y \in G$$

and at least one of them is \mathfrak{M} -measurable, then there exist a continuous additive function $A: G \rightarrow H$ and constants $a, b \in H$ such that

$$\begin{cases} f(x) - A(x) - a \in K, \\ g(x) - A(x) - b \in K, \\ h(x) - A(x) - a + b \in K \end{cases}$$

for $x \in G$.

Acknowledgement. The research was supported by the Silesian University Mathematics Department (Iterative Functional Equations and Real Analysis program).

References

- [1] Baron K., Kannappan Pl., *On the Pexider difference*, Fund. Math. **134** (1990), 247–254.
- [2] Baron K., Volkman P., *On orthogonally additive functions*, Publ. Math. Debrecen **52** (1998), 291–297.
- [3] Brzdęk J., *On orthogonally exponential functionals*, Pacific J. Math. **181** (1997), 247–267.

- [4] Kochanek T., Wyrobek-Kochanek W., *Measurable orthogonally additive functions modulo a discrete subgroup*, Acta Math. Hungar. **123** (2009), 239–248.
- [5] Rätz J., *On orthogonally additive mappings*, Aequationes Math. **28** (1985), 35–49.
- [6] Sikorska J., *On a Pexiderized conditional exponential functional equation*, Acta Math. Hungar. **125** (2009), 287–299.
- [7] Wyrobek W., *Orthogonally additive functions modulo a discrete subgroup*, Aequationes Math. **78** (2009), 63–69.

INSTITUTE OF MATHEMATICS
SILESIAN UNIVERSITY
BANKOWA 14
40-007 KATOWICE
POLAND
e-mail: wwyrobek@math.us.edu.pl