

AN ALTERNATIVE EQUATION FOR GENERALIZED POLYNOMIALS OF DEGREE TWO

ZOLTÁN BOROS , RAYENE MENZER 

Abstract. In this paper we consider a generalized polynomial $f: \mathbb{R} \rightarrow \mathbb{R}$ of degree two that satisfies the additional equation $f(x)f(y) = 0$ for the pairs $(x, y) \in D$, where $D \subseteq \mathbb{R}^2$ is given by some algebraic condition. In the particular cases when there exists a positive rational m fulfilling

$$D = \{ (x, y) \in \mathbb{R}^2 \mid x^2 - my^2 = 1 \},$$

we prove that $f(x) = 0$ for all $x \in \mathbb{R}$.

1. Introduction

Let \mathbb{R} , \mathbb{Q} , and \mathbb{N} denote the set of all real numbers, rationals, and positive integers, respectively.

We call a function $f: \mathbb{R} \rightarrow \mathbb{R}$ *additive* if

$$f(x + y) = f(x) + f(y)$$

for all $x, y \in \mathbb{R}$. The function f is called *\mathbb{Q} -homogeneous* if the equation $f(qx) = qf(x)$ is fulfilled by every $q \in \mathbb{Q}$ and $x \in \mathbb{R}$. As it is well-known

Received: 25.04.2023. Accepted: 03.10.2023. Published online: 26.10.2023.

(2020) Mathematics Subject Classification: 39B55, 39B72.

Key words and phrases: additive functions, quadratic functions, generalized polynomials, alternative equation.

Research of Z. Boros has been supported by the K-134191 NKFIH Grant. Project no. K134191 have been implemented with the support provided from the National Research, Development and Innovation Fund of Hungary, financed under the K_20 funding scheme.

©2023 The Author(s).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License CC BY (<http://creativecommons.org/licenses/by/4.0/>).

(see M. Kuczma [6, Theorem 5.2.1]), if $f: \mathbb{R} \rightarrow \mathbb{R}$ is additive, then f is \mathbb{Q} -homogeneous as well. For more information concerning these notions the reader is referred to the monograph [6].

A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is called *quadratic* if it satisfies the Jordan-von Neumann functional equation:

$$f(x+y) + f(x-y) = 2f(x) + 2f(y)$$

for all $x, y \in \mathbb{R}$. In what follows we will apply the fact that f is quadratic if and only if there exists a bi-additive and symmetric functional $B: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = B(x, x)$ for every $x \in \mathbb{R}$ (see e.g. J. Aczél, J. Dhombres [1, Chapter 11, Proposition 1]). Quadratic functions are also called generalized monomials of degree 2. Further, additive functions are generalized monomials of degree 1 and real constants are generalized monomials of degree 0. Generalized polynomials are defined as sums of generalized monomials of respective degrees. For more facts on generalized polynomials the reader is referred to [6, Chapter 15.9] and L. Székelyhidi [8]. In particular, we call a function $f: \mathbb{R} \rightarrow \mathbb{R}$ *generalized polynomial of degree two* if we can write it with the following decomposition $f(x) = g(x) + a(x) + b$ for every $x \in \mathbb{R}$, where $g: \mathbb{R} \rightarrow \mathbb{R}$ is a quadratic function, $a: \mathbb{R} \rightarrow \mathbb{R}$ is additive and $b \in \mathbb{R}$. Here we do not exclude the particular cases when $g(x) = 0$ or $a(x) = 0$ identically, or $b = 0$. Therefore, we consider constant functions and additive mappings (as well as their sums) as particular generalized polynomials of degree two.

For any positive rational m we define the following sets:

$$S_0 = \{(x, y) \in \mathbb{R}^2 \mid xy = 1\},$$

$$S_{1,m} = \{(x, y) \in \mathbb{R}^2 \mid x^2 - my^2 = 1\},$$

$$S_2 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}.$$

Z. Kominek, L. Reich and J. Schwaiger [5] investigated additive real functions that satisfy the additional equation

$$(1) \quad f(x)f(y) = 0$$

for every $(x, y) \in D$, considering various subsets D of \mathbb{R}^2 . In several cases they obtained $f(x) = 0$ for every $x \in \mathbb{R}$. Their result for $D = S_2$ was extended by Z. Boros and W. Fechner [2] to the situation when f is a generalized polynomial. On the other hand, P. Kutas [7] has recently established the existence of a non-zero additive function $f: \mathbb{R} \rightarrow \mathbb{R}$ fulfilling (1) for all $(x, y) \in S_0$. The case of bounded $f(x)f(y)$ on S_2 was investigated by these authors [3] for particular generalized polynomials of degree two. In a recent paper [4] the present authors obtained analogous results, including one for $D = S_{1,m}$,

assuming that f is a generalized monomial. However, it remained an open problem whether one can extend the latter result to the more general case when f is a generalized polynomial of an arbitrary degree. The purpose of the present paper is to prove such a theorem when f is a generalized polynomial of degree two.

2. Main results

Now we can establish our main theorems.

THEOREM 2.1. *Let m denote a positive rational. Suppose that $f: \mathbb{R} \rightarrow \mathbb{R}$ is a generalized polynomial of degree two and $f(x)f(y) = 0$ for all solutions of the equation $x^2 - my^2 = 1$. Then f is identically equal to zero.*

PROOF. Given a generalized polynomial f of degree two, we can associate a quadratic function $g: \mathbb{R} \rightarrow \mathbb{R}$, an additive function $a: \mathbb{R} \rightarrow \mathbb{R}$ and a constant $b \in \mathbb{R}$ with f such that

$$(2) \quad f(x) = g(x) + a(x) + b$$

for every $x \in \mathbb{R}$. Moreover, there exists a symmetric and bi-additive functional $G: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ such that $g(x) = G(x, x)$ for every $x \in \mathbb{R}$.

Now, let $x \in \mathbb{R}$ be such that $x \geq 1$. Then there exists $0 \leq y \in \mathbb{R}$ such that $x^2 - my^2 = 1$. If α, β are rational numbers such that $\alpha^2 - m\beta^2 = 1$, it is easy to verify the equality

$$(\alpha x + \beta my)^2 - m(\beta x + \alpha y)^2 = 1,$$

hence our assumptions on f imply

$$(3) \quad f(\alpha x + \beta my)f(\beta x + \alpha y) = 0.$$

Considering the decomposition (2) of f we can calculate that

$$\begin{aligned} f(\alpha x + \beta my) &= g(\alpha x + \beta my) + a(\alpha x + \beta my) + b \\ &= \alpha^2 g(x) + (\beta m)^2 g(y) + 2\alpha\beta m G(x, y) + \alpha a(x) + \beta m a(y) + b \end{aligned}$$

and

$$\begin{aligned} f(\beta x + \alpha y) &= g(\beta x + \alpha y) + a(\beta x + \alpha y) + b \\ &= \beta^2 g(x) + \alpha^2 g(y) + 2\alpha\beta G(x, y) + \beta a(x) + \alpha a(y) + b. \end{aligned}$$

Due to equation (3), for every pair of rationals (α, β) fulfilling $\alpha^2 - m\beta^2 = 1$, at least one of the foregoing expressions is equal to zero.

What is more, we can find infinitely many distinct pairs (α_j, β_j) such that $\alpha_j^2 - m\beta_j^2 = 1$ and both α_j and β_j are rationals. Namely, let

$$\alpha_j = \frac{mj^2 + 1}{mj^2 - 1} \quad \text{and} \quad \beta_j = \frac{2j}{mj^2 - 1}$$

for $j \in \mathbb{N}$ such that $mj^2 \neq 1$.

Thus, for every $j \in N_m \doteq \mathbb{N} \setminus \{1/\sqrt{m}\}$, we have either

$$\begin{aligned} 0 &= \left(\frac{mj^2 + 1}{mj^2 - 1}\right)^2 g(x) + \left(\frac{2jm}{mj^2 - 1}\right)^2 g(y) + \frac{4jm(mj^2 + 1)}{(mj^2 - 1)^2} G(x, y) \\ &\quad + \frac{mj^2 + 1}{mj^2 - 1} a(x) + \frac{2jm}{mj^2 - 1} a(y) + b \end{aligned}$$

or

$$\begin{aligned} 0 &= \left(\frac{2j}{mj^2 - 1}\right)^2 g(x) + \left(\frac{mj^2 + 1}{mj^2 - 1}\right)^2 g(y) + \frac{4j(mj^2 + 1)}{(mj^2 - 1)^2} G(x, y) \\ &\quad + \frac{mj^2 + 1}{mj^2 - 1} a(y) + \frac{2j}{mj^2 - 1} a(x) + b. \end{aligned}$$

Multiplying both equations by $(mj^2 - 1)^2$ and introducing the functions

$$\begin{aligned} P(j) &= j^4 m^2 (g(x) + a(x) + b) + j^3 m^2 (4G(x, y) + 2a(y)) \\ &\quad + j^2 (2mg(x) + 4m^2 g(y) - 2mb) \\ &\quad + j(4mG(x, y) - 2ma(y)) + g(x) - a(x) + b \end{aligned}$$

and

$$\begin{aligned} \tilde{P}(j) &= j^4 m^2 (g(y) + a(y) + b) + j^3 m (4G(x, y) + 2a(x)) \\ &\quad + j^2 (4g(x) + 2mg(y) - 2mb) \\ &\quad + j(4G(x, y) - 2a(x)) + g(y) - a(y) + b, \end{aligned}$$

we have $P(j) = 0$ or $\tilde{P}(j) = 0$ for each integer $j \in N_m$. Hence either P or \tilde{P} has infinitely many zeros. On the other hand, both P and \tilde{P} are polynomials of degree not greater than 4. Therefore, one of them has to be identically

equal to 0. So either each coefficient of P equals zero, or each coefficient of \tilde{P} equals zero. In the first case, considering the coefficient of j^4 , we obtain

$$m^2(g(x) + a(x) + b) = 0,$$

which obviously implies

$$f(x) = g(x) + a(x) + b = 0.$$

Now let us consider the second case, when we obtain the following system of equations:

$$(4) \quad g(y) + a(y) + b = 0,$$

$$(5) \quad 4G(x, y) + 2a(x) = 0,$$

$$(6) \quad 4g(x) + 2mg(y) - 2mb = 0,$$

$$(7) \quad 4G(x, y) - 2a(x) = 0,$$

$$(8) \quad g(y) - a(y) + b = 0.$$

By summing (4) and (8) we get $g(y) = -b$. Then substituting the value of $g(y)$ in (6) we get $g(x) = mb$. Finally, by summing (5) and (7) we get $G(x, y) = 0$ and $a(x) = 0$.

We have thus proved, for an arbitrary real number $x \geq 1$, that either $f(x) = 0$ or we have $a(x) = 0$ and $g(x) = mb$.

Suppose that $f(x_0) \neq 0$ for some $x_0 \geq 1$. Then we must have $a(x_0) = 0$, $g(x_0) = mb$ and

$$0 \neq f(x_0) = g(x_0) + a(x_0) + b = mb + 0 + b = (m + 1)b,$$

which implies $b \neq 0$. Then we also have $2x_0 \geq 2$ (and thus $2x_0 \geq 1$),

$$g(2x_0) = 4g(x_0) = 4mb \neq mb$$

and

$$\begin{aligned} f(2x_0) &= g(2x_0) + a(2x_0) + b = 4g(x_0) + 2a(x_0) + b \\ &= 4mb + 0 + b = (4m + 1)b \neq 0 \end{aligned}$$

since $b \neq 0$ and $m \neq -\frac{1}{4}$ (as m is positive). So our conclusions for $2x_0$ do not satisfy our previous results for all $x \geq 1$. This contradiction shows that such an element x_0 does not exist.

We have thus proved $f(x) = 0$ for every real number $x \geq 1$. Applying [4, Lemma 2.1], we obtain that $f(x) = 0$ for all $x \in \mathbb{R}$. \square

COROLLARY 2.2. *Let a and b denote positive real numbers such that $\frac{a^2}{b^2}$ is rational. Suppose that $f: \mathbb{R} \rightarrow \mathbb{R}$ is a generalized polynomial of degree two and $f(x)f(y) = 0$ for all solutions of the equation $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$. Then f is identically equal to zero.*

PROOF. Let u and w be real numbers fulfilling the condition $u^2 - \frac{a^2}{b^2}w^2 = 1$. Moreover, let $f_a(t) = f(at)$ for all $t \in \mathbb{R}$. Clearly, then f_a is a generalized polynomial of degree two as well. For $x = au$ and $y = aw$ we have

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = u^2 - \frac{a^2}{b^2}w^2 = 1,$$

hence our assumption yields

$$f_a(u)f_a(w) = f(au)f(aw) = f(x)f(y) = 0.$$

Therefore f_a satisfies the assumptions in Theorem 2.1 with $m = \frac{a^2}{b^2}$, hence f_a is identically equal to zero, which yields $f(x) = f_a(x/a) = 0$ for every $x \in \mathbb{R}$ as well. \square

References

- [1] J. Aczél and J. Dhombres, *Functional Equations in Several Variables*, Encyclopedia Math. Appl., **31**, Cambridge University Press, Cambridge–New York–New Rochelle–Melbourne–Sydney, 1989.
- [2] Z. Boros and W. Fechner, *An alternative equation for polynomial functions*, Aequationes Math. **89** (2015), no. 1, 17–22.
- [3] Z. Boros, W. Fechner, and P. Kutas, *A regularity condition for quadratic functions involving the unit circle*, Publ. Math. Debrecen **89** (2016), no. 3, 297–306.
- [4] Z. Boros and R. Menzer, *An alternative equation for generalized monomials*, Aequationes Math. **97** (2023), no. 1, 113–120. DOI: 10.1007/s00010-022-00917-y
- [5] Z. Kominek, L. Reich, and J. Schwaiger, *On additive functions fulfilling some additional condition*, Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II **207** (1998), 35–42.
- [6] M. Kuczma, *An Introduction to the Theory of Functional Equations and Inequalities*, second edition, Birkhäuser Verlag, Basel–Boston–Berlin, 2009.
- [7] P. Kutas, *Algebraic conditions for additive functions over the reals and over finite fields*, Aequationes Math. **92** (2018), no. 3, 563–575.
- [8] L. Székelyhidi, *Regularity properties of polynomials on groups*, Acta Math. Hungar. **45** (1985), no. 1-2, 15–19.

ZOLTÁN BOROS
INSTITUTE OF MATHEMATICS
UNIVERSITY OF DEBRECEN
P. O. Box: 400
H-4002 DEBRECEN
HUNGARY
e-mail: zboros@science.unideb.hu

RAYENE MENZER
INSTITUTE OF MATHEMATICS
UNIVERSITY OF DEBRECEN
P. O. Box: 400
H-4002 DEBRECEN
HUNGARY
e-mail: rayene.menzer@science.unideb.hu, riene1997@gmail.com